

Effects of changing spatial resolution on the results of landscape pattern analysis using spatial autocorrelation indices

Ye Qi¹ and Jianguo Wu^{2,*}

¹*Scripps Institution of Oceanography, University of California at San Diego, La Jolla, CA 92093-0220, USA;* ²*Biological Sciences Center, Desert Research Institute, University of Nevada System, P.O. Box 60220, Reno, Nevada 89506-0220, USA*

Keywords: landscape patterns, spatial analysis, spatial autocorrelation, scale effect, grain size, Moran Coefficient, Geary Ratio, Cliff-Ord statistic

Abstract

Understanding the relationship between pattern and scale is a central issue in landscape ecology. Pattern analysis is necessarily a critical step to achieve this understanding. Pattern and scale are inseparable in theory and in reality. Pattern occurs on different scales, and scale affects pattern to be observed. The objective of our study is to investigate how changing scale might affect the results of landscape pattern analysis using three commonly adopted spatial autocorrelation indices, *i.e.*, Moran Coefficient, Geary Ratio, and Cliff-Ord statistic. The data sets used in this study are spatially referenced digital data sets of topography and biomass in 1972 of Peninsular Malaysia. Our results show that all three autocorrelation indices were scale-dependent. In other words, the degree of spatial autocorrelation measured by these indices vary with the spatial scale on which analysis was performed. While all the data sets show a positive spatial autocorrelation across a range of scales, Moran coefficient and Cliff-Ord statistic decrease and Geary Ratio increases with increasing grain size, indicating an overall decline in the degree of spatial autocorrelation with scale. The effect of changing scale varies in their magnitude and rate of change when different types of landscape data are used. We have also explored why this could happen by examining the formulation of the Moran coefficient. The pattern of change in spatial autocorrelation with scale exhibits threshold behavior, *i.e.*, scale effects fade away after certain spatial scales are reached (for elevation). We recommend that multiple methods be used for pattern analysis whenever feasible, and that scale effects must be taken into account in all spatial analysis.

Introduction

Landscapes are mosaics of patches that differ in size, shape, and contents (Risser *et al.* 1984; Forman and Godron 1986; Wu and Levin 1994). Numerous studies have shown that the spatial pattern of landscapes may have significant influences on ecological processes, such as population dynamics, biogeochemical cycling, and aspects of biodiversity

(*e.g.*, Burgess and Sharpe 1981; Zonneveld and Forman 1990; Opdam 1991; Wiens *et al.* 1993; Wu *et al.* 1993; Wu and Vankat 1991a, b, 1995; Wu and Levin 1994). Therefore, identifying and characterizing spatial pattern across a range of scales (especially large ones) using various quantitative methods are often necessary in landscape ecological studies (Turner and Gardner 1991; Cullinan and Thomas 1992; Wu 1992a).

* To whom correspondence should be addressed: Biological Sciences Center, Desert Research Institute, University of Nevada System, P.O. Box 60220, Reno, Nevada 89506-0220, USA. Email: jwu@maxey.unr.edu

Data and methods

The two data sets used for this study are georeferenced digital data of elevation and above-ground biomass in Peninsular Malaysia (Fig. 1a, b; see Brown *et al.* 1994 for more details). Information in these data sets is represented using a rectangular lattice or grid, each cell of which has one or more values. Both data sets have 220 rows and 188 columns, with a grid cell size (the minimum or basic spatial unit, BSU) of 2.25×2.25 kilometers.

To change the grain size across a range of spatial scales, we systematically aggregated the data from the original spatial resolution to larger areal aggregates in the following way. Each BSU was treated as one basic unit, and therefore the grain size at this scale was expressed as 1 by 1. A 2×2 areal unit, then, corresponded to the grain size that contained four BSUs (two on each side). This was accomplished by aggregating four adjacent basic areal units, assigning the arithmetic mean of the four to the newly formed areal unit. This procedure was repeated until the entire region of the data sets was covered. In total, 20 different grain sizes (spatial scales) were created, ranging from 1×1 through 20×20 BSUs (*i.e.*, $1, 2^2, 3^2, \dots, 20^2$). Table 1 gives detailed information on the grain sizes, the number of areal units at grain size, and the starting position of aggregation at each scale.

In the aggregation process, sometimes the original data sets had to be modified (edge rows or columns were omitted or repeated) to obtain integer numbers of rows and columns. For example, in some cases we trimmed off the first 8 columns because they consisted mainly of zeros, representing ocean or areas of no concern. In some other cases, more rows were needed to maintain an integral number of rows for moving up to the next spatial scale, and we duplicated the first and/or last row. While this kind of modification was necessary only for technical convenience, we believed that this modification would not affect the results of our analysis because of the relatively large size of the dates.

Moran Coefficient (MC), Geary Ratio (GR) and Cliff-Ord statistic (CO) are among the most commonly used indices for the analysis of spatial auto-

correlation in geographically referenced ecological data (Cliff and Ord 1973, 1981; Odland 1988; Goodchild 1986; Griffith 1988; Legendre 1993). These autocorrelation coefficients are defined as follows (Griffith 1988).

(1) Moran Coefficient

$$MC = \left(n / \sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij} \right) \sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij} (x_i - \bar{x})(x_j - \bar{x}) / \sum_{i=1}^{i=n} (x_i - \bar{x})^2 \quad (1)$$

(2) Geary Ratio

$$GR = \left((n-1) / (2 \sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij}) \right) \sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij} (x_i - x_j)^2 / \sum_{i=1}^{i=n} (x_i - \bar{x})^2 \quad (2)$$

(3) Cliff-Ord statistic

$$CO = \left(n / \sum_{i=1}^{i=n} \sum_{j=1}^{j=n} w_{ij} \right) \sum_{i=1}^{i=n} \sum_{j=1}^{j=n} w_{ij} (x_i - \bar{x})(x_j - \bar{x}) / \sum_{i=1}^{i=n} (x_i - \bar{x})^2 \quad (3)$$

$$\text{with } \sum_{i=1}^{i=n} w_{ij} = 1 \text{ and } 0 \leq w_{ij} \leq 1 \quad (4)$$

where n is the total number of areal units over the entire landscape, x_i and x_j are values of areal units i and j , and \bar{x} is the mean of all areal units. c_{ij} denotes the connectivity between areal units i and j , and it takes a value of 1 if areal units i and j are adjacent and 0 otherwise). $C = \{c_{ij}\}$ is called the connectivity matrix ($n \times n$). w_{ij} is the standardized connectivity between i and j , (*i.e.*, $w_{ij} = c_{ij} / \sum_j c_{ij}$), and $W = \{w_{ij}\}$ is the standardized connectivity matrix (Griffith 1988).

Two areal units i and j are considered adjacent if they are within certain distance from each other. In practice, the values of c_{ij} and hence w_{ij} are determined simply by checking if two areal units i and j are immediately next to each other. In other words,




Fig. 1. The landscape data sets of Peninsular Malaysia: (a) elevation, where the actual altitude increases with the degree of darkness with category 0 denoting the sea level; (b) above-ground biomass in 1972, where areas with greater biomass are represented by darker pixels. All the data sets have 220 rows and 188 columns, and have a spatial resolution of 2.25×2.25 kilometers.

areal units i and j are regarded as being adjacent if they share a line. In our case where a regular rectangular lattice is used, a grid cell can only be adjacent to the neighboring cells on the four sides (*i.e.*, upper, lower, left, and right).

All three statistics have the following common properties: (1) there exists an expected value (mean)

that corresponds to the case of no spatial autocorrelation, (2) the coefficient is able to discriminate between different spatial arrangements of the same set of values $\{x_i\}$ on a two-dimensional surface, and (3) all of them also have well established distributions based on conventional sampling theory (Griffith 1988; Odland 1988). The spatial correla-




Fig. 1b.

tion indices, as defined above, are determined by two factors: the value of each areal unit and the spatial relationship among all the areal units. In all our calculations, the spatial extent was fixed, so that our study was focused completely on the scale effects from a grain size point of view. We calculated the three spatial autocorrelation indices at each of the 20 spatial scales that were specified by different grain sizes for the two data sets. The three

indices were used together for comparing their sensitivity to scale change and verifying the algorithm we developed. We realized that altering grain sizes would involve two closely related but different issues: change in the area of the spatial unit at different grain sizes and arrangement or configuration of spatial units at smaller scales to form higher level aggregates (e.g., Wu *et al.* 1994). However, this study was concentrated only on the first issue.

Table 1. Layout of the aggregating scheme from the basic spatial unit (1×1 BSU) to larger areal unit aggregates (up to 20×20 BSUs). Grain size, starting row, and starting column are measured or denoted by the basic spatial units, while the numbers of rows and columns represent the actual numbers of the areal units corresponding to each spatial scales (grain sizes or levels of aggregation).

Grain size	#Areal units at each scale	Starting row	Starting column
1 x 1	220 x 180	3	9
2 x 2	110 x 90	3	9
3 x 3	73 x 60	4	9
4 x 4	55 x 45	3	9
5 x 5	44 x 36	3	9
6 x 6	36 x 30	4	9
7 x 7	31 x 26	4	7
8 x 8	27 x 23	4	5
9 x 9	24 x 20	4	9
10 x 10	22 x 18	3	9
11 x 11	20 x 16	3	11
12 x 12	18 x 15	4	9
13 x 13	17 x 14	3	7
14 x 14	15 x 13	5	7
15 x 15	14 x 12	6	9
16 x 16	13 x 11	7	11
17 x 17	13 x 11	3	2
18 x 18	12 x 10	4	9
19 x 19	11 x 9	7	11
20 x 20	11 x 9	3	9

Constructing the connectivity matrix is usually the most critical step for calculating spatial autocorrelation indices (Griffith 1988). It is often the computational bottleneck that limits the size of the spatial data sets (Griffith 1990). There are, in total, 41,360 grid cells in our original data set (220 rows by 188 columns). This would produce a connectivity matrix of 41,360 by 41,360, which is too large for a personal computer to handle. Fortunately, the property of a regular rectangular lattice of data allows us to reduce the computational demand by changing the way of dealing with the connectivity matrix. In general, for a spatial data set with k rows and p columns, there are at most $4kp-2k-2p$ non-zero values in the connectivity matrix of $kp \times kp$. Therefore, we can use a matrix of $4 \times kp$, instead of $kp \times kp$, to record all the information contained in the connectivity matrix. By so doing, the computer memory usage can be reduced by a factor of




Fig. 2. Scale effect on spatial autocorrelation indices (Moran Coefficient – MC, Geary Ratio – GR, and Cliff-Ord statistic – CO): (a) elevation, and (b) biomass.

$kp/4$, as compared to the conventional algorithm (e.g., Griffith 1990). In this study, k and p are 220 and 188, so the computational demand is cut down by a factor of 10,340 (i.e., $4 \times 188 \times 220 = 165,440$ elements in the connectivity matrix in total).

Results and analysis

For the three autocorrelation indices used here, the values for both Moran coefficient and Cliff-Ord statistic usually fall in between -1 and 1, though they may exceed 1.0 for some weights and attribute values (Goodchild 1986). The theoretical value for no spatial autocorrelation for these two indices is $-1/(n-1)$. A value of less than zero indicates a nega-

tive spatial autocorrelation, and a value larger than zero suggests a positive spatial autocorrelation. The value of Geary Ratio, on the other hand, ranges from 0 to 2, with 1 for no, larger than 1 for negative, and smaller than 1 for positive spatial autocorrelation. Therefore, a positive spatial autocorrelation is always detected if the computed value falls between 0 and 1, no matter which of the three autocorrelation coefficients is used.

We shall discuss the results of our analysis by addressing the following three related questions. (1) Based on the two landscape data sets, how does changing grain size affect the results of analysis using the three spatial autocorrelation correlation coefficients? (2) For a given spatial autocorrelation index, how do such scale-dependent changes vary with different types of landscape data? (3) Why do such differences occur for various types of landscape data?

To explore the first question above, we examined the sensitivity of the three autocorrelation coefficients to changes in spatial resolution by applying them simultaneously to the same data set. Figure 2 shows how the numerical values of these coefficients respond to increasing grain sizes. In all the analyses with the two data sets, the computed values for Moran coefficient and Cliff-Ord statistic are very closely related across a wide range of scales. This may be expected from a scrutiny of the similar formulation of the two indices (see Eqs. 1 and 3). For the elevation data set, Moran coefficient and Cliff-Ord statistic decrease steadily from changing grain size affect the results of analysis using the three spatial autocorrelation correlation coefficients? (2) For a given spatial autocorrelation index, how do such scale-dependent changes vary 0.96 to about 0.68, as the grain size increases from the minimum spatial unit to the 11 by 11 BSU aggregate, and then tend to level off (Fig. 2a; Table 2). Geary ratio, on the other hand, increases almost linearly first from 0.038 to 0.37 as the grain size increases to the scale of 11 by 11 basic areal units, and then gradually levels off around the value of 0.4 (Fig. 2a; Table 2).

For the biomass data set, Moran coefficient and Cliff-Ord statistic again decline with increasing grain size (Fig. 2b). The decrease continues all the

Table 2. Values of the three spatial autocorrelation coefficients for different data sets with changing grain size. The grain size is denoted by the basic spatial areal units (BSUs) for these data sets.

Grain size	Elevation			Biomass		
	MC	GR	CO	MC	GR	CO
1	0.9691	0.0382	0.9620	0.7647	0.2314	0.7691
2	0.9370	0.0764	0.9237	0.7468	0.2607	0.7416
3	0.8998	0.1180	0.8834	0.7336	0.2698	0.7339
4	0.8638	0.1609	0.8433	0.7227	0.2794	0.7237
5	0.8332	0.1961	0.8090	0.7246	0.2750	0.7242
6	0.8076	0.2259	0.7837	0.7333	0.2687	0.7360
7	0.7751	0.2630	0.7489	0.7173	0.2836	0.7202
8	0.7435	0.2999	0.7169	0.7190	0.2759	0.7272
9	0.7398	0.3092	0.7087	0.7318	0.2657	0.7324
10	0.6948	0.3558	0.6677	0.6997	0.2963	0.7064
11	0.6822	0.3695	0.6532	0.6862	0.3057	0.6900
12	0.6727	0.3891	0.6432	0.6763	0.3288	0.6779
13	0.7083	0.3454	0.6732	0.6671	0.3372	0.6679
14	0.6703	0.3961	0.6353	0.6750	0.3338	0.6732
15	0.6700	0.3947	0.6319	0.6669	0.3262	0.6605
16	0.6646	0.3948	0.6251	0.6294	0.3836	0.6271
17	0.6558	0.4095	0.6198	0.6174	0.3721	0.6146
18	0.6375	0.4428	0.5941	0.6054	0.4158	0.5859
19	0.6689	0.4140	0.6124	0.5839	0.4406	0.5603
20	0.6620	0.4217	0.6032	0.5644	0.4615	0.5453

way to the grain size of 20 by 20 BSUs, but it seems more gradual (from 0.76 to 0.54 for the entire range of scales, see Table 2). Geary ratio shows a reversed trend with increasing spatial scale as it does for the elevation data set. All three indices show a rather distinctive change (through) around the grain size of 11 by 11 basic areal units in the data sets of elevation. We speculate that this may correspond to an abrupt change in spatial patchiness of some sort around that scale in the peninsular Malaysia landscape. However, we cannot completely rule out the possibility that it may be a result of some artifact in the data aggregation procedure. This phenomenon deserves further examination in future analysis of this type, especially when data sets of higher spatial resolution and larger extent are available.

Given the existence of obvious effects of changing scale on the results of spatial autocorrelation coefficients as demonstrated above, one of the questions in order is whether there is any generality about them. In this particular case study, we explore how scale effects manifest themselves with the different types of landscape data sets under con-




Fig. 3. Comparison of scale effects on Moran Coefficient among different landscape data sets.

Table 3. Values of Moran coefficient computed at the smallest, middle and largest spatial scales for the two different data sets. The changes in Moran coefficient between any two of the three grain sizes are also given in the table, with the percentage changes in the parentheses.

Data type	MC value			Δ MC		
	1×1	10×10	20×20	$1 \times 1 - 20 \times 20$	$1 \times 1 - 10 \times 10$	$10 \times 10 - 20 \times 20$
Elevation	0.9691	0.6948	0.6620	0.3071	0.2743 (89.3%)	0.0328 (10.7%)
Biomass	0.7647	0.6997	0.5644	0.2003	0.0650 (32.5%)	0.1353 (67.5%)

sideration. The answer to this question has already been touched upon earlier in the discussion of the results that show how the three indices are affected by changing scales using the same data sets. To further elucidate this question, however, we choose to use Moran coefficient to highlight the comparison (Fig. 3). The choice of Moran coefficient is based on the observation that the three indices used in this study are in good agreement, with Moran coefficient and Cliff-Ord statistic exhibiting great similarity in both pattern of change and numerical values.

From Fig. 3, some differences in the pattern of change in spatial autocorrelation with scale are appreciable. There is a noticeable contrast in the amplitude of decrease in the degree of autocorrelation from the smallest spatial unit (1×1 BSU) to the

largest areal aggregate (20×20 BSUs) for the two data sets, with the maximum decrease of 0.3071 for elevation and a minimum of 0.2003 for biomass. In addition, the rate of decrease (*i.e.*, the steepness of each curve) is different for these data sets. For instance, the degree of spatial autocorrelation for biomass data drops appreciably more slowly than for elevation. Also, the biomass data set shows a faster changing pace over the larger spatial scales (from 10×10 BSU to 20×20 BSU areal aggregates) than smaller ones (from 1×1 BSU to 10×10 BSU aggregates), which is in contrast with elevation (Table 3).

Why does Moran coefficient show different scale effects for different types of landscape data sets? One way to explore this question is to examine the formulation of the autocorrelation index. The




Fig. 4. Effect of changing spatial scale on Moran Coefficient and its two aggregated terms: (a) elevation, and (b) biomass.

formula used to calculate Moran coefficient may

be decomposed into four terms: n , $\sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij}$, $\sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij}(x_i - \bar{x})(x_j - \bar{x})$, and $\sum_{i=1}^{i=n} (x_i - \bar{x})^2$ (see

Eq. 3). It does not seem to make any sense to analyze scale effects using each term separately. However, when we regroup them into two terms, $n / \sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij}$ and $\sum_{i=1}^{i=n} \sum_{j=1}^{j=n} c_{ij} (x_i - \bar{x})(x_j - \bar{x}) / \sum_{i=1}^{i=n} (x_i - \bar{x})^2$, we are able to gain insights into the question being addressed here. With increasing grain size, the first term exhibits a nearly linear increase, and the second term decreases which essentially resembles the behavior of Moran coefficient as a whole (Fig. 4a–b). Apparently, the decreasing trend

of the second term overwhelms the increasing trend of the first term and, therefore, the overall pattern of change in spatial autocorrelation is primarily determined by that of the first term. A similar dissection of the formulas for Cliff-Ord statistic and Geary ratio can also be done, and the outcome is expected to be similar. The first term in the Moran coefficient formula is affected only by spatial aggregation or partitioning of the original data set, while the second term is influenced by both the manipulation of grain size and the numerical value of each grid cell. Therefore, the first term has the same response to all the data types of the same partitioning, whereas the second term presents different responses for different landscape data sets due to their numerical differences.

Discussion and conclusions

It is common in landscape ecology to represent spatial data with a regularly divided rectangular lattice that is composed of a large number of equal-sized grid cells. The dimension of the grid cell determines the spatial resolution or grain size of the data set. Such geographically referenced data sets of different spatial resolutions are often used and spatial aggregation is frequently needed in landscape ecological studies. For instance, to assess and monitor the change in biomass in a region, one may use SPOT, Thematic Mapper, MSS, or AVHRR satellite data, each of which has a different spatial resolution (16×16 meters for SPOT, 30×30 meters for TM, 180×180 meters for MSS, and 1×1 kilometers for AVHRR). To advance our understanding of spatial processes in landscape ecology, therefore, it is critically important and absolutely imperative to first understand how changing spatial scale (e.g., grain size) affects the results of spatial analysis of landscape patterns. Our study on spatial autocorrelation coefficients sheds some light on this issue.

Our study shows that both data sets (elevation and biomass) of Peninsula Malaysia are positively spatially autocorrelated across a range of scales (from 2.25×2.25 km to 45×45 km). From the results discussed earlier, it is clear that changing

spatial scale significantly affects the values of all three autocorrelation indices. Therefore, spatial analysis of landscape pattern using these indices at single scales, in general, may provide little useful, or even misleading information. Specifically, as the grain size increases, the values of Moran coefficient and Cliff-Ord statistic decrease, while Geary ratio increases. In other words, the degree of spatial autocorrelation as measured by these three indices, in general, decreases with increasing spatial scale. This result is not readily intuitive. It appears that for certain types of landscape data (*e.g.*, elevation), there exists a spatial threshold beyond which scale effects are no longer obvious. Yet, for other types of landscape data (*e.g.*, biomass) such a threshold does not seem to occur or, if it does, it would be on a much larger spatial scale that is beyond the range limit considered in our study. Although some geographical studies suggested that these three indices could behave differently (*e.g.*, Cliff and Ord 1973, 1981), we found no appreciable difference among them with regularly gridded data sets used in this study.

Given the scale effects, the results of all spatial analysis should be presented with explicit specification of the scale on which the study is conducted. Whenever feasible, an examination of scale effects on the analytical results across a range of scales that are relevant to the landscape pattern under investigation is most desirable. In the case of spatial autocorrelation analysis, the three indices do not seem to differentiate significantly from each other in terms of the ability to detect the scale effects according to our study. Yet, we concur with other researchers that multiple methods should be used wherever possible for the sake of comparison and verification (Turner *et al.* 1989; Cullinan and Thomas 1992). Now that landscape patterns and methods to analyze them are both scale dependent, an important task for landscape ecologists today is to develop not only techniques to detect characteristic spatial domains (Wiens and Milne 1989), but also the ways to scale up and down pattern and process in various landscapes (Turner, Dale and Gardner 1989; Constanza and Maxwell 1994; Wu and Levin 1994).

Acknowledgements

We would like to thank Sandra Brown and Louis Iverson for providing the data sets. We are also grateful to Robert Burgess, Arthur Getis, Daniel Griffith, Charles Hall, Louis Iverson, and Gil Pontius for reviewing the manuscript. During this study, Y. Qi was financially supported through a research grant from the U.S. Department of Energy awarded to Charles Hall. Financial support for J. Wu was in part from the National Science Foundation under Cooperative Agreement OSR-9353227.

References

- Allen, T.F.H. and Starr, T.B. 1982. *Hierarchy: Perspectives for Ecological Complexity*. University of Chicago Press, Chicago.
- Burgess, R.L. and Sharpe, D.M. (eds). 1981. *Forest Island Dynamics in Man-Dominated Landscapes*. Springer-Verlag, New York.
- Brown, S., Iverson, L.R. and Lugo, A.E. 1994. Land use and biomass changes of forests in Peninsular Malaysia during 1972–1982: use of GIS analysis. *In* *Effects of Land-Use Change on Atmospheric CO₂ Concentrations: Southeast Asia as a Case Study*. pp. 117–143. Edited by V. Dale. Springer-Verlag, New York.
- Cliff, A. and Ord, J.K. 1973. *Spatial Autocorrelation*. Pion, London.
- Cliff, A. and Ord, J.K. 1981. *Spatial Process*. Pion, London.
- Costanza, R. and Maxwell, T. 1994. Resolution and predictability: An approach to the scaling problem. *Landsc. Ecol.* 9: 47–57.
- Cullinan, V.I. and Thomas, J.M. 1992. A comparison of quantitative methods for examining landscape pattern and scale. *Landsc. Ecol.* 7: 211–227.
- Forman, R.T.T. and Godron, M. 1986. *Landscape ecology*. Wiley, New York.
- Fortin, M.-J., Drapeau, P. and Legendre, P. 1989. Spatial autocorrelation and sampling design in plant ecology. *Vegetatio* 83: 209–222.
- Getis, A. and Boots, B. 1978. *Models of spatial processes: an approach to the study of point, line and area patterns*. Cambridge University Press, Cambridge, England.
- Getis, A. and Ord, J.K. 1992. The analysis of spatial association by use of distance statistics. *Geographical Analysis* 24: 189–206.
- Goodchild, M.F. 1986. *Spatial Autocorrelation. Concepts and Techniques in Modern Geography* 47, Geo Books, Norwich.
- Greig-Smith, P. 1952. The use of random and contiguous quadrats in the study of the structure of plant communities. *Ann. Bot., New Series* 16: 293–316.
- Griffith, D.A. 1988. *Advanced Spatial Statistics. Special Topics*

- in the Exploration of Quantitative Spatial Data Series. Kluwer Academic Publishers, Dordrecht.
- Griffith, D.A. 1990. Supercomputing and spatial statistics: A reconnaissance. *Prof. Geogr.* 42(4): 481–492.
- Kershaw, K.A. 1957. The use of cover and frequency in the detection of pattern in plant communities. *Ecology* 38: 291–299.
- Legendre, P. 1993. Spatial autocorrelation: trouble or new paradigm? *Ecology* 74: 1659–1673.
- Legendre, P. and Fortin, M.J. 1989. Spatial pattern and ecological analysis. *Vegetatio* 80: 107–138.
- Levin, S.A. 1992. The problem of pattern and scale in ecology. *Ecology* 73: 1943–1967.
- Levin, S.A., Powell, T. and Steele, J.H. (eds.) 1993. *Patch Dynamics*. Springer-Verlag, New York.
- Meentemeyer, V. and Box, E.O. 1987. Scale effects in landscape studies. *In* *Landscape Heterogeneity and Disturbance*. pp. 15–34. Edited by M.G. Turner. Springer-Verlag, New York.
- Milne, B.T. 1988. Measuring the fractal geometry of landscapes. *Appl. Math. Comp.* 27: 67–79.
- Morris, D.W. 1987. Ecological scale and habitat use. *Ecology* 68: 362–369.
- Nellis, M.D. and Briggs, J.M. 1989. The effect of spatial scale on Konza landscape classification using textural analysis. *Landscape Ecol.* 3: 93–100.
- Odland, J. *Spatial Autocorrelation*. SAGE Publications, Newbury Park.
- O'Neill, R.V., DeAngelis, D.L., Waide, J.B. and Allen, T.F.H. 1986. *A Hierarchical Concept of Ecosystems*. Princeton University Press, Princeton.
- O'Neill, R.V., Krummel, J.R., Gardner, R.H., Sugihara, G., Jackson, B., DeAngelis, D.L., Milne, B.T., Turner, M.G., Zygmunt, B., Chistensen, S., Dale, V.H. and Graham, R.L. 1988. Indices of landscape pattern. *Landscape Ecol.* 1: 153–162.
- Opdam, P. 1991. Metapopulation theory and habitat fragmentation: A review of holarctic breeding bird studies. *Landscape Ecol.* 5: 93–106.
- Risser, P.G., Karr, J.R. and Forman, R.T.T. 1984. *Landscape Ecology: Directions and Approaches*. Illinois Natural History Survey, Special Publication Number 2.
- Turner, M.G., Dale, V.H. and Gardner, R.H. 1989. Predicting across scales: theory development and testing. *Landscape Ecol.* 3: 245–252.
- Turner, M.G. and Gardner, R.H. (eds). 1991. *Quantitative Methods in Landscape Ecology*. Springer-Verlag, New York.
- Turner, M.G., O'Neill, R.V., Gardner, R.H. and Milne, B.T. 1989. Effects of changing spatial scale on the analysis of landscape pattern. *Landscape Ecol.* 3: 153–162.
- Wiens, J.A. and Milne, B.T. 1989. Scaling of “landscape” in landscape ecology, or, landscape ecology from a beetle's perspective. *Landscape Ecol.* 3: 87–96.
- Wiens, J.A., Stenseth, N.C., Horne, B. van and Ims, R.A. 1993. Ecological mechanisms and landscape ecology. *Oikos* 66: 369–380.
- Wu, J. 1992a. Detecting spatial patterns: the net-function interpolation. *Coenoses* 7(3): 137–143.
- Wu, J. 1992b. Balance of nature and environmental protection: A paradigm shift. *Proc. 4th Intern. Conf. Asia Experts*. Portland State University, Portland, OR, May 7–8, 1992.
- Wu, J., Jelinski, D. and Qi, Y. 1994. Spatial pattern analysis of a boreal forest landscape: Scale effects and interpretation. *Proc. VI Intern. Conf. Ecology (INTECOL)*, pp. 165. Univ. of Manchester, Manchester, August 21–26, 1994.
- Wu, J. and Levin, S.A. 1994. A spatial patch dynamic modeling approach to pattern and process in an annual grassland. *Ecol. Monogr.* 64(4): 447–464.
- Wu, J., Li, B. and Wu, Y. 1992. Patchiness and patch dynamics: I. Concepts and mechanisms. *Chin. J. Ecol.* 11(4): 41–45.
- Wu, J. and Vankat, J.L. 1991a. A system dynamics model of island biogeography. *Bull. Math. Biol.* 53: 911–930.
- Wu, J. and Vankat, J.L. 1991b. An area-based model of species richness dynamics of forest islands. *Ecol. Modell.* 58: 249–271.
- Wu, J. and Vankat, J.L. 1995. Island biogeography: Theory and applications. *In* *Encyclopedia of Environmental Biology*. Edited by W.A. Nierenberg: Academic Press (in press).
- Wu, J., Vankat, J.L. and Barlas, B. 1993. Effects of patch connectivity and arrangement on animal metapopulation dynamics: a simulation study. *Ecol. Modell.* 65: 221–254.
- Zonneveld, I.S. and Forman, R.T.T. (eds). 1990. *Changing Landscapes: An Ecological Perspective*. Springer-Verlag, New York.